

Lectade Concentrate Oral Rehydration Therapy for Dogs and Cats

A ready to use electrolyte replacement for scouring dogs and cats.

Product Information

Active Constituents

- Glucose | undiluted 278.75 g/L | diluted (1 part to 11.5 parts) 22.30 g/L
- Glycine | undiluted 38.56 g/L | diluted (1 part to 11.5 parts) 3.08 g/L
- Citric Acid Monohydrate | undiluted 3.28 g/L | diluted (1 part to 11.5 parts) 0.26 g/L
- Potassium Citrate | undiluted 0.75 g/L | diluted (1 part to 11.5 parts) 0.06 g/L
- Potassium Phosphate Monobasic | undiluted 25.53 g/L | diluted (1 part to 11.5 parts) 2.04 g/L
- Sodium chloride | undiluted 53.44 g/L | diluted (1 part to 11.5 parts) 4.28 g/L

Indications

- Diarrhoea
- Dehydration
- Post-surgical or disease convalescence
- Concomitant treatment of viral or bacterial scours



Lectade Concentrate Oral Rehydration Therapy Sachets

A water soluble electrolyte replacer for scouring animals.

Product Information

Active Constituents

- Glucose | undiluted - 44.61 g | diluted as "Lectade Solution" - 22.30 g/L
- Glycine | undiluted - 6.18 g | diluted as "Lectade Solution" - 3.08 g/L
- Citric Acid Monohydrate | undiluted - 0.48 g | diluted as "Lectade Solution" - 0.26 g/L
- Potassium Citrate | undiluted - 0.12 g | diluted as "Lectade Solution" - 0.06 g/L
- Potassium Phosphate Monobasic | undiluted - 4.08 g | diluted as "Lectade Solution" - 2.04 g/L
- Sodium chloride | undiluted - 8.58 g | diluted as "Lectade Solution" - 4.28 g/L

Indications

- Diarrhoea
- Dehydration (simple or associated with diseases such as canine parvovirus and feline panleucopaenia)
- Post travelling stress and dehydration
- Post-surgical or disease convalescence
- Concomitant treatment of viral or bacterial scours

